

Australian hepatic types at the National Herbarium of New South Wales (NSW) and the Ray Herbarium (SYD)

E.A. Brown, H.P. Ramsay and J. Seur

Abstract

E.A. Brown, Ramsay, H.P. & Seur, J. (National Herbarium of New South Wales, Royal Botanic Gardens, Sydney, NSW, Australia 2000) 1992. Australian hepatic types at the National Herbarium of New South Wales (NSW) and the Ray Herbarium (SYD). Telopea 5(1): 67–90. The National Herbarium of New South Wales and the Ray Herbarium in Sydney, New South Wales hold important collections of type specimens of Australian hepatics. A total of 99 names are listed here, together with details of the protologue, complete label data for each specimen, other herbaria known to hold part of the type, and recent information concerning the status of the name. Included are many Stephani and Carrington and Pearson names as well as more recent ones described by Hewson and Na-Thalang.

Introduction

While the Hepatic Collections in the two herbaria in Sydney, New South Wales, are not large [6000 specimens, 100 types (NSW) and 2284 specimens, <10 types (SYD)] (Ramsay & Seur 1990), type specimens of many Australian and exotic species are present. Particularly significant collections held at the National Herbarium of New South Wales [NSW] are those of T. Whitelegge, W.W. Watts (including some from W. Gunn), and H.J. Hewson; and at the Ray Herbarium, University of Sydney [SYD], those of Hewson and O. Na-Thalang. The main geographical regions represented by the types are Australia including Lord Howe Island (Whitelegge, Watts, Na-Thalang), presented in this paper, and Vanuatu [New Hebrides] (W. Gunn) and Papua New Guinea (Hewson) to be reported separately.

The earliest descriptions of New South Wales hepatics are those of B. Carrington & W.H. Pearson (1888) who determined and published a list of Whitelegge's collections. R.H. Bastow's work on Tasmanian hepatics (1888, 1914) also included reference to many New South Wales collections. Watts published several papers on Australian hepatics (1902, 1903, 1905) and left a manuscript (1912), as part of a Census of New South Wales Plants, that he had intended publishing in the same way as he had done for the mosses (Watts & Whitelegge 1902, 1906). In this unpublished manuscript he listed species, localities and collector's names for 205 Hepatics and 10 Anthocerotales. The names contained in this unpublished manuscript include many described by F. Stephani based on specimens sent to him for identification (Stephani 1884, 1889; Stephani & Watts 1914).

The cryptogamic collection at NSW was increased greatly in the late 19th and early 20th century by W.W. Watts. After his departure to Victoria in 1916 (Ramsay 1980) curation and addition to the collections was very limited. Conditions at NSW were extremely cramped until 1982 when a new building opened and the W.W. Watts room was set aside for cryptogams.

A period of inactivity in collecting and describing hepatics in New South Wales persisted from 1920 until the mid 1960s, when taxonomic research was undertaken

by G.K. Berrie and R. Carolin's students Hewson and Na-Thalang, at Sydney University. In 1963 Berrie published *Haplomitrium intermedium*, the first member of the Calobryales recorded in Australia, from a type collected in the Blue Mountains (Berrie 1963). Hewson (1970*a, b*) revised the family Aneuraceae in Australia and New Guinea while Na-Thalang (1980) revised the genus *Riccia* in Australia. Hewson and Na-Thalang types are located either at SYD or NSW and in some cases at both.

There is still much to be learnt about hepatics in New South Wales; we do not even have a complete species list for this State in which both subtropical and temperate taxa occur. Some recent publications (Ramsay *et al.* 1990, Downing *et al.* 1991) and further studies being undertaken by H.P. Ramsay *et al.* list the hepatics for specific areas in N.S.W. and form the basis for the eventual production of a Census of New South Wales hepatics.

Locating types is essential for taxonomic revisions and a register providing details and locations of type specimens can be useful. Ramsay & Seur (1990) and Ramsay *et al.* (1990) outlined the methods for collection and recording of data for a register of type collections of mosses. Their investigations highlighted the often incomplete information in many type citations in taxonomic literature (e.g. date of collection and collector's number). Since most descriptions of species of hepatics published prior to the 1970s do not cite the herbarium holding the type the holotype cannot be clearly defined. Examining the type specimen and its label data may give important details not included in the protologue.

Method

In this paper we deal only with those hepatic type specimens of Australian species held in the two herbaria named NSW and SYD.

The studies were carried out in a series of steps:

1. All data on each herbarium packet containing specimens designated as types or sp. nov. (i.e. putative types) were recorded on index cards.
2. The type description for each taxon was obtained and the label data compared with that in the protologue. Where these matched, the specimen is recorded as a probable type.
3. Where possible, reference is made to the herbarium holding the major collections of the author of the basionym if this is not NSW or SYD. In most cases, except for species named after the mid 1970s, the holotypes are held in European herbaria and specimens in Australia usually are isotypes. For a number of species a series of specimen numbers were listed with the type description and are syntypes and it may be desirable, therefore, when taxonomic revisions are carried out, to designate a lectotype at that time.
4. References are given for recent studies, name changes, lectotypification etc. for each taxon.
5. To ensure that we have detected all relevant type specimens held at NSW and SYD, we checked publications and searched through all collections at NSW and SYD for specimens not labelled as types but carrying the original label data. As the status of quite a number of specimens had been overlooked in the repackaging at NSW during the early 1980s these further checks, although time-consuming, proved rewarding.
6. The information is now stored on disk and will remain available for future updating.

The register of type specimens of Australian hepatics presented here has been arranged alphabetically by genera and species based on the name for which the specimen is a type. The list contains the original name for each taxon and reference, the protologue [Proto] (collection details as cited in the protologue), full label data [Label] for each separate specimen (label 1, 2 etc.) at NSW and/or SYD if more than one was present. Also listed are location and status of all types [Types] where known (from examination or from literature), synonymy where applicable, as well as any other recent information such as lectotypification [Comment].

At some time in the past many of Watts' collections have been broken up, repacketed and relabelled; the duplicates may all be contained within the one specimen folder or sometimes they have been separated. Where label information does not differ significantly between packets it is usually listed only once, with a note that there are 2 or more specimens; sometimes information on packets within the same specimen folder differs markedly and this has been recorded. The labels on the outer specimen folders has not been recorded as it is only a more recent copy of the information on the Watts' packets.

No taxonomic decisions other than those already in the literature are presented here, nor is lectotypification of any species implied. This is particularly important for taxa described by Stephani; in his later years he broke up many of his specimens and sold duplicates to BM, FH and other institutions. Some of these duplicates, when they come from mixed collections, are not the taxon referred to in the protologue and great care must be taken when typifying. It is probably unwise and inaccurate, therefore, to cite holotypes and this has not been done, even when Stephani specified a collector's number. Herbaria, other than NSW and SYD, holding types have been listed from *Index Hepaticarum*. *Index Hepaticarum* VIII (Bonner 1976) and prior volumes were used to obtain putative label information for many of the types held at G. E.A. Brown has examined the type specimens from overseas herbaria for the genera *Riccardia* and *Aneura* in the course of other studies (Brown & Braggins 1989).

Register of names

Aneura aequicellularis Stephani (Stephani & Watts 1914: 95)

PROTO: 'Australia, New South Wales (Wentworth Falls): Watts legit No. 117 [1117].'

LABEL: NEW SOUTH WALES: Wentworth Falls, cave, W.W. W[atts] 1117, Sep 1912.

TYPES: G (lectotype), NSW (isolectotype).

COMMENT: = *Riccardia aequicellularis* (Stephani) Hewson fide Hewson (1970b: 79).
Lectotypified by Brown & Braggins (1989: 26).

Aneura athertonensis Hewson (1970a: 188)

PROTO: 'Holotype — Charmillan Creek, Tully Falls Road, Atherton Tableland, on basalt rock on creek bank in disturbed rain-forest, Qld. Hewson, 422, 8.1964 (NSW): Isotype (BRI).'

LABEL: QUEENSLAND: Charmillan Creek, rock, rainforest, basalt, H.J. Hewson H.64.71.422, 22 Aug 1964.

TYPES: NSW (179239: holotype), BRI (isotype).

COMMENT: Name in current use.

***Aneura coriacea* Stephani (1917: 23)**

PROTO: 'Australia: New South Wales (Watts legit.)'

LABEL: NEW SOUTH WALES: Katoomba Falls, W.W. *Watts* 567, 15 Jan 1902.

TYPES: G (11036: lectotype), NSW (H241: isolectotype).

COMMENT: No particular specimen cited in protologue. The NSW specimen has been annotated by Hewson: 'Isotype of *A. coriacea* = *Riccardia crassa*, published Hewson 1970b'. Lectotypified by Brown & Braggins (1989: 85). Specimen in G seen by Stephani is *Watts* 567 (Icones Hep. 474).

***Aneura eachamensis* Hewson (1970a: 184)**

PROTO: 'Holotype — Vision Falls, Lake Eacham, Atherton Tableland, on basalt rock in rain-forest, Queensland. Hewson, 398, 8.1964 (NSW): Isotype (BRI).'

LABEL: QUEENSLAND: Vision Falls, Lake Eacham, H.J. *Hewson* H.64.68.398, 20 Aug 1964.

TYPES: NSW (179240: holotype), BRI (isotype).

COMMENT: Name in current use.

***Aneura gigantea* Stephani (Stephani & Watts 1914: 95)**

PROTO: 'Australia, N.S. Wales, (Cambewarra): Watts legit No. 920.'

LABEL: NEW SOUTH WALES: under cliff, above Baldy's, Cambewarra, W.W. *Watts* 920, 18 Oct 1907 [3 packets in one specimen folder].

TYPES: G (11042), NSW.

COMMENT: = *Aneura alterniloba* (Hook. f. & Taylor) Taylor & Hook. f. in Gottsche, Lindenb. & Nees var. *gigantea* (Stephani) Hewson; G 11042 and '920 NSW' are cited as the holotype and isotype respectively by Hewson (1970a: 187).

***Aneura pusilla* Stephani (Stephani & Watts 1914: 96)**

PROTO: 'Australia, (Old Railway Cutting, Blackheath): Watts legit, 1051.'

LABEL: NEW SOUTH WALES: old railway cutting, Blackheath, W.W. *Watts* 1051, Jan 1911 [3 packets in one specimen folder].

TYPES: G (11049: lectotype), NSW (H242: isolectotype).

COMMENT: Listed as a synonym of *Riccardia crassa* (Schwaegr.) Carrington & Pearson (Hewson 1970a: 104). Taxon reinstated and transferred to *Riccardia*; lectotypified by Brown & Braggins (1989: 82).

***Aneura rodwayi* Hewson (1970a: 188)**

PROTO: 'Holotype—Hartz River, Mt. Hartz National Park, on clay soil near river bank, Tas., Hewson, 215, 9.1963, (NSW).'

LABEL: TASMANIA: Hartz River, Mt Hartz, on soil near river, H.J. *Hewson* H63.33.215, 10 Sep 1963.

TYPE: NSW (179241: holotype).

COMMENT: Name in current use.

***Aneura rufescens* Stephani (Stephani & Watts 1914: 96)**

PROTO: 'Australia, New South Wales, (National Pass, Wentworth Falls): Watts legit, 1124.'

LABEL: NEW SOUTH WALES: National Pass, Wentworth Falls, W.W. *Watts* 1124, Sep 1912 [2 packets in one specimen folder].

TYPES: G (11050: lectotype), NSW (H243: isolectotype).

COMMENT: = *Riccardia crassa* fide Hewson (1970b: 104). Lectotypified by Brown & Braggins (1989: 85).

Aneura rupicola Stephani (1917: 41)

PROTO: 'Australia: New South Wales (Watts legit).'

LABEL: NEW SOUTH WALES: 1. Shaws Bay, log, W.W. *Watts* 579, 20 Sep 1901; 2. Shaws Bay, Ballina, R[ichmond] R[iver], log, W.W. *Watts* 579, 20 Sep 1901 [both packets are in one specimen folder].

TYPES: G (11051 ex Hb Levier 3388), NSW (H248).

COMMENT: G specimen annotated 'Original. Shaw's Bay, east Ballina. Icones Hep. 520'. = *Riccardia rupicola* (Stephani) Hewson; G 11051 cited as holotype and 'NSW 579' as isotype by Hewson (1970b: 99).

Aneura walesiana Stephani (Stephani & Watts 1914: 97)

PROTO: 'Australia, New South Wales, (Blackheath): Watts legit, 1023.'

LABEL: NEW SOUTH WALES: Horseshoe Falls, Blackheath, wet cliff, W.W. *Watts* 1023, 7 Jan 1911 [3 packets in one specimen folder].

TYPES: G (lectotype), NSW (H244: isolectotype).

COMMENT: = *Riccardia crassa* fide Hewson (1970b: 104). Lectotypified by Brown & Braggins (1989: 85).

Aneura wattiana Stephani (1917: 46)

PROTO: 'Australia, New South Wales (Watts legit).'

LABEL: NEW SOUTH WALES: 1. E Ballina, [Richmond River], on tree fern, wet places, W.W. *Watts* 237, 2 Mar 1901 (NSW H252); 2. label identical, no NSW number.

TYPES: G (11057 ex Hb Levier 3018), NSW.

COMMENT: No specific locality or specimen number cited in protologue. Specimen in G (11057) has the following data 'W.W. *Watts* 237, 1901, Ballina, New South Wales. Icones Hep. 299'. = *Riccardia wattiana* (Stephani) Hewson; G 11057 cited as holotype and 'NSW 237' as isotype by Hewson (1970b: 91). The un-numbered NSW specimen has a second Watts collection attached to it (*Watts* 346).

Archilejeunea australis Stephani (1911: 734)

PROTO: 'Australia, New South Wales.'

LABEL: NEW SOUTH WALES: back of Wickham's, nr. Ballina, tree, W.W. *Watts* 415, 3 June 1902.

TYPES: G (ex Hb Levier 3483), NSW.

COMMENT: = *Thysananthus australis* (Stephani) Thiers & Gradstein; specimen in G (ex Hb Levier 3483-*Watts* 415) cited as the holotype by Thiers & Gradstein (1989: 66); no reference is made to the G specimen being labelled as the type by Stephani.

Balantiopsis decurrens *Stephani* (Stephani & Watts 1914: 98)

PROTO: 'Australia, (Wyong): Watts, 985.'

LABEL: NEW SOUTH WALES: gully south of house, 'Kingwell', Wyong, W.W. *Watts* 985, Nov 1907 [3 packets in one specimen folder].

TYPES: G, NSW.

COMMENT: = *B. diplophylla* (Hook. f. & Taylor) Mitten fide Engel (1968: 109, 111).

Balantiopsis hastatistipula *Stephani* (Stephani & Watts 1914: 98)

PROTO: 'Australia, (Blackheath), Watts, 1052.'

LABEL: NEW SOUTH WALES: Blackheath, old railway cutting, W.W. *Watts* 1052, Jan 1911 [2 packets in one specimen folder].

TYPES: G, NSW.

COMMENT: = *Isotachis* sp. fide Engel (1968: 118).

Balantiopsis kingwella *Stephani* (Stephani & Watts 1914: 99)

PROTO: 'Australia (Kingwell, Wyong): Watts 943.'

LABEL: NEW SOUTH WALES: gully south of house, 'Kingwell', Wyong, W.W. *Watts* 943, Nov 1907 [3 packets in one specimen folder].

TYPES: G, NSW.

COMMENT: = *B. diplophylla* fide Engel (1968: 109, 111).

Balantiopsis subkingwella *Stephani* (Stephani & Watts 1914: 99)

PROTO: 'Australia (Kingwell, Wyong): Watts 945, 971^a.'

LABEL: NEW SOUTH WALES: 1. 'Kingwell', Wyong, W.W. *Watts* 945, 20 Aug 1909; 2. 'Kingwell', Wyong, gully south of house, W.W. *Watts* 971^a, Nov 1907 (NSW H285) [2 packets in one specimen folder].

TYPES: G, NSW.

COMMENT: = *B. diplophylla* fide Engel (1968: 109, 111). *Watts* 971^b is the type of *Mastigobryum corbieri* Stephani, see below.

Brachiolejeunea robusta *Stephani* (1912: 141)

PROTO: 'Australia. New South Wales.'

LABEL: NEW SOUTH WALES: 1. Hopkins Creek, Stanwell Park, rocks, W.W. *Watts* 830, May 1905; 2. Hopkins Creek, Stanwell Park, [W.W.] *Watts* 830, May 1905.

TYPES: FH, G (ex Hb Levier 4882), NSW, U.

COMMENT: Specimen in G (Hb Levier 4882-*Watts* 830) cited as holotype and FH as isotype by Thiers & Gradstein (1989: 64) but there is no supporting discussion. Reduced to synonymy under *Spruceanthus thozetianus* (Gottsche & F. Muell.) Thiers & Gradstein.

Chiloscyphus maximus *Stephani* (Stephani & Watts 1914: 103)

PROTO: 'Australia (Etta's Glen, Black Spur, Vict.): Watts, 968.'

LABEL: VICTORIA: Etta's Glen, Black's Spur, Healesville, W.W. *Watts* 968 & 968^a, 21 Dec 1906 [5 packets in one specimen folder].

TYPES: G, NSW.

COMMENT: The NSW specimen consists of 5 small packets variously labelled *Watts 968*, *Watts 968a* or *Watts 968 + 968a*. All are annotated '*Chiloscyphus maximus* St. sp. n. 1913' in Watts' hand. Apparently a large variety of *C. coalitus* (Hook.) Nees in Gottsche, Lindenb. & Nees (Scott 1985: 133).

Chiloscyphus montanus Stephani (Stephani & Watts 1914: 103)

PROTO: 'Australia (Neate's Glen, Blackheath): Watts, 927.'

LABEL: NEW SOUTH WALES: Rotunda, Neate's Glen, B[lack]heath, on damp ground base of cliff, W.W. Watts 927, 7 Apr 1903 [3 packets in one folder].

TYPES: G, NSW.

COMMENT: = *Geocalyx caledonicus* Stephani fide Grolle (1979).

Fimbriaria dioica Stephani (Stephani & Watts 1914: 104)

PROTO: 'Australia, N.S. Wales (near Gladesville): Watts 1095.'

LABEL: NEW SOUTH WALES: off Pittwater Road, Gladesville, W.W. Watts 1095 & [J.F.] Shirley, Nov 1911 [3 specimens in one folder].

TYPES: G, NSW.

COMMENT: *Fimbriaria* is an illegitimate name but this taxon has not been formally transferred to *Asterella*, where it belongs. This group is under revision.

Fossombronina grossepapillata Stephani (1917: 73)

PROTO: 'Australia (Watts legit).'

LABEL: NEW SOUTH WALES: park, Young, (mining holes), W.W. Watts 988, Sep-Oct 1906 [3 packets in one folder].

TYPES: G (22171), NSW.

COMMENT: G 22171 (*Watts 988*) designated as type specimen by Scott & Pike (1988: 196).

Fossombronina integrifolia Stephani (1917: 73)

PROTO: 'Australia orientalis (N.S. Wales).'

LABEL: NEW SOUTH WALES: 1. Parramatta, E. Cheel (*Hb Watts 799*), 1 Oct 1910 [2 packets]; 2. Rookwood, Sydney, E. Cheel (*Hb Watts 812*), 5 Aug 1900 [2 packets].

TYPES: FH (812), G (22174), NSW.

COMMENT: Specimens of *Watts 812* held at FH and G designated as the type by Scott & Pike (1988: 197).

Fossombronina wattsii Stephani (1917: 75)

PROTO: 'Australia, New South Wales (Watts legit).'

LABEL: NEW SOUTH WALES: The Gap, near Young, W.W. [Watts] 881, 26 Sep 1905.

TYPES: G (*Hb Levier 5155*), NSW.

COMMENT: No locality or specimen number cited in protologue but G specimen of *Watts 881* cited as type in Bonner (1965: 221). Type material (at NSW) checked and studied by Pike: 'abnormal spores present 8/3/82'.

Frullania asperifolia Stephani (Stephani & Watts 1914: 105)

PROTO: 'Australia (Yarrangobilly Caves & Mount Wilson): Watts 1093, etc).'

LABEL: NEW SOUTH WALES: 1. Yarrangobilly Caves, W.W. *Watts* 923, Jan 1906; 2. 'Yarrawa', Mt Wilson, W.W. *Watts* 1063, 10 Jan 1911; 3. Denmarque, Mt Wilson, W.W. *Watts* 1078, 13 Jan 1911; 4. Falls beyond Kiandra road, Yarrangobilly Caves, W.W. *Watts* 926, Jan 1906; 5. Yarrangobilly [Caves], gully beyond hostel, W.W. *Watts* 922, Jan 1906; 6. Avenue, Mt Wilson, W.W. *Watts* 1093, 11 Jan 1911.

TYPES: G (18193), NSW.

COMMENT: Although two localities were cited in the protologue only one number (*Watts* 1093) was given and this is cited as the type by Bonner (1965: 239). Hattori (1979a) cites the type as 'G18193, the Avenue, Mt Wilson, 3500 ft on tree Jan 11, 1911, leg. W.W. Watts (Hep. Aust. ex Herb. Watts 109 [= 1093])' and considers this taxon conspecific with *F. falciloba* Taylor ex Lehm.

Frullania australis Stephani (1911: 659)

PROTO: 'Australia orientalis, New South Wales.'

LABEL: NEW SOUTH WALES: E Ballina, tree, W.W. *Watts* 117, 14 Aug 1900 [3 packets in one folder].

TYPES: G (18194 ex Hb Levier 2638), NSW.

COMMENT: G 18194 is cited as the type by Bonner (1965: 241), without the Watts collection number. Known only from the type collection (Hattori 1979b: 123).

Frullania belmorensis Stephani (Stephani & Watts 1914: 106)

PROTO: 'Australia (Belmore Falls): Watts, 931.'

LABEL: NEW SOUTH WALES: Belmore Falls, W.W. *Watts* 931, 5 Oct 1908 [3 packets in one folder].

TYPES: G, NSW.

COMMENT: Known only from the type collection (Hattori 1979b: 126).

Frullania difficilis Stephani (1889: 160)

PROTO: 'Australia, New South Wales.'

LABEL: NEW SOUTH WALES: Park, Gray's Island, Richmond River, W.W. *Watts* 125, 6 May 1900.

TYPES: G (ex Hb Levier 2632), MEL 61558, NSW.

COMMENT: No specimen number or specific locality is cited in the protologue but the type is cited by Bonner (1965: 288) as 'New South Wales, Richmond River, Pearce's Creek, 2 VIII 1900, W.W. Watts s.n., Hb Levier 2632 in Hb G'. Hattori (1987: 553) lists a number of other possible syntypes (including the above NSW specimen) and places *F. difficilis* in synonymy under *Frullania (Trachycolea) seriata* Gottsche ex Stephani.

Frullania elongata Stephani (1910: 423)

PROTO: 'Australia orientalis.'

LABEL: NEW SOUTH WALES: Pearce's Creek nr R. Trimble's, bark of old tree stump, W.W. *Watts* 96, 2 Aug 1900.

TYPES: G, NSW.

COMMENT: *Watts* 96 is cited by Bonner (1965: 296) as the type. = *F. rubella* fide Hattori (1979a: 354); = *F. rubella* Gottsche ex Stephani var. *elongata* (Stephani) Hattori fide Hattori (1983: 165).

Frullania excisula Stephani (Stephani & Watts 1914: 107)

PROTO: 'Australia, N.S. Wales (Mount Wilson and Blackheath): *Watts* 1030, 1071.'

LABEL: NEW SOUTH WALES: 1. Horseshoe Falls to Jungle, Blackheath, (rock by track), *W.W. Watts* 1030, 7 Jan 1911 [2 packets]; 2. Yarrowa, Mt Wilson, *W.W. Watts* 1071, 11 Jan 1911 [3 packets].

TYPES: G, MEL, NSW.

COMMENT: Bonner (1965: 229) cited *Watts* 1017 [= 1071] as the type. The NSW specimens of both *Watts* 1030 and 1071 are annotated by S. Hattori Apr. 1986: 'not type of *F. squarrulosa* but syntype of *F. excisula*'. *F. excisula* seems to be conspecific with *Frullania falciloba* (Hattori 1979a: 336).

Frullania ferdinandi-muelleri Stephani (1910: 417)

PROTO: 'Australia, New South Wales.'

LABEL: NEW SOUTH WALES: Tintenbar(?), R[ichmond] R[iver], tea tree bark, *W.W. Watts* 153, 2 Aug 1900.

TYPES: G (18211 ex Hb Levier 2628), NSW.

COMMENT: No information on locality, date, collector or specimen number(s) appears in the protologue but Hattori (1979a) describes this species as known only from the type collection and cites G 18211 as the type.

Frullania filipendula Stephani (1910: 419)

PROTO: 'Australia, Queensland, New South Wales.'

LABEL: NEW SOUTH WALES: E Ballina, base of saplings and on ground, heath, *W.W. Watts* 468, 17 June 1902 [2 packets in folder].

TYPES: G (18212 ex Hb Levier 3416), NSW.

COMMENT: No information on locality, date, collector or specimen number(s) appears in the protologue but Bonner (1965: 304) cites this specimen (G 18212) as the type. Seems to be conspecific with *F. rubella* (Hattori 1979a: 340).

Frullania howeana Stephani (Stephani & Watts 1914: 107)

PROTO: 'Lord Howe Island. (*Watts* 62).'

LABEL: NEW SOUTH WALES: near Face, east side of Mt Lidgbird, L[ord] H[owe] I[land], *W.W. Watts* 62, 27 July 1911 [2 packets in folder].

TYPES: G (18215), NSW.

COMMENT: Endemic, known only from the type collection (Hattori 1979a: 346).

Frullania minutistipula Stephani (Stephani & Watts 1914: 108)

PROTO: 'Australia (Rodriguez Pass): *Watts* 1001^b, 1001^c.'

LABEL: NEW SOUTH WALES: Rodriguez Pass, Blackheath, *W.W. Watts* 1001^b & *c*, 4 Jan 1911.

TYPES: G, NSW.

COMMENT: NSW specimen has been annotated (by Watts?): 'mixed with *Frullania gracillima* sp. n. and *Radula wattsiiana*'. = *F. rostrata* (Hook. f. & Taylor) Hook. f. & Taylor ex Gottsche, Lindenb. & Nees fide Hattori (1979a: 352).

Frullania simmondsii Stephani (Stephani & Watts 1914: 109)

PROTO: 'Australia (near Brisbane): Simmonds leg. (Watts 1110).'

LABEL: QUEENSLAND: near Brisbane, *J.H. Simmonds, Hb Watts 1110, 27 Aug 1887.*

TYPES: G, NSW.

COMMENT: NSW specimen annotated as isotype by Hattori. Known only from the type collection (Hattori 1979b: 143).

Frullania subtropica Stephani (1910: 416)

PROTO: 'Australia, New South Wales.'

LABEL: NEW SOUTH WALES: off Tintenbar road, 3 miles [5 km] from Ballina, Richmond River, tree, *W.W. Watts 107, 30 June 1900.*

TYPES: G (18230 ex Hb Levier 2602), NSW.

COMMENT: Protologue does not include locality, date, collector or specimen information but G 18230 is cited as the type by Bonner (1965: 440).

Frullania variabilis Stephani (1910: 420)

PROTO: 'Australia, New South Wales.'

LABEL: NEW SOUTH WALES: Cambewarra Mtn, *W.W. [Watts] 628, 22 May 1903.*

TYPES: G (ex Hb Levier 4416), NSW.

COMMENT: Protologue does not include locality, date, collector or specimen information but the above specimen in G (Hb Levier 4416) is cited as the type by Bonner (1965: 462). Known only from the type collection; probably a form of *F. pycnantha* (Hook. f. & Taylor) Taylor ex Gottsche, Lindenb. & Nees (Hattori 1979b: 149).

Haplomitrium intermedium Berrie (1963: 191)

PROTO: 'Between one and two miles along the road from Bilpin to Mount Irvine, Blue Mountains, N.S.W. HOLOTYPE: Sydney University Herbarium, Bryophyte No. 2/62. COTYPE: Professor R.M. Schuster Collection No. 50,616.'

LABEL: see below

TYPES: SYD (holotype; not located).

COMMENT: Probably on loan, but record not found.

Isotachis grandis Carrington & Pearson (1888: 1041)

PROTO: 'On wet rocks, Lawson, Blue Mountains, June, 1884.'

LABEL: NEW SOUTH WALES: 1. Lawson, Blue Mountains, on wet rocks, [*T. Whitelegge*], June 1884; 2. Lawson, no 14 liverwort, [*T. Whitelegge*], June 1884.

TYPES: BM, MANCH?, NSW.

COMMENT: The former NSW specimen has been further annotated by Whitelegge: '17 *Isotachis grandis* n. sp. (small portion of stem) pLXXVI 34 TW W.H.P.' and the latter has the additional information: '17 N. sp. No 34 P. *Isotachis grandis*'. Probably no. 34 of a batch sent to Pearson by Whitelegge.

Isotachis terricola Stephani (Stephani & Watts 1914: 10)

PROTO: 'Australia (Blackheath): Watts, 1016.'

LABEL: NEW SOUTH WALES: short cut to Neate's Glen, Blackheath, creek, swamp, W.W. *Watts 1016*, 5 Jan 1911 [3 packets in folder].

TYPES: G, NSW.

COMMENT: = *I. intortifolia* (Hook. f. & Taylor) Gottsche fide Hatcher (1960: 602).

***Jungermannia wattiana Stephani* (1917: 96)**

PROTO: 'Australia, New South Wales (Watts legit.)'

LABEL: NEW SOUTH WALES: 1. Alst[onville] Cutting, 7 miles [11 km] nearly, W.W. *Watts 303*, 24 June 1901; 2. Alstonville Cutting, R[ichmond] R[iver], nearly 7 miles [11 km] from Ballina, W.W. *Watts 303*, 24 June 1901.

TYPES: G (14307 ex Hb Levier 3141), NSW.

COMMENT: No specimen numbers given in protologue but Bonner (1976: 410) cites the type as: 'NSW, Alstourith [= Alstonville] cutting, 7 miles from Ballina, 24.June.1901, Rev. Watts 303, Hb Levier 3141, Hb G 14307.'

***Jungermannia whiteleggei Carrington & Pearson* (1888: 1051)**

PROTO: no specimens cited.

LABEL: NEW SOUTH WALES: Moore Park, [Sydney], [*T. Whitelegge*] 37, Dec 1884.

TYPES: K or MANCH?, NSW.

COMMENT: The NSW specimen has been annotated by Whitelegge: '*Jungermannia whiteleggei* n. sp., pl XXXII 36 T.W. W.H.P.' It is probably no. 36 of a batch sent to W.H. Pearson. No specimen information is given in the protologue but the information on the packet is consistent with that found on other type specimens sent by Whitelegge to Pearson. = *Chaetophyllopsis whiteleggei* (Carrington & Pearson) R. Schuster fide Schuster (1960: 69).

***Lejeunea wildii Stephani* (1889: 165)**

PROTO: 'Queensland, leg. C. Wild. - Herb. Brotherus.'

LABEL: QUEENSLAND: Hamilton, C.[J.] Wild, July 1887 [2 identically labelled specimens].

TYPES: G (15847), BM, MANCH, MEL, NSW (H495 & H496).

COMMENT: G 15847 cited as holotype and BM, MANCH and MEL as isotypes by Thiers & Gradstein (1989: 6); almost certainly the only specimen matching the protologue. Reduced to synonymy under *Acrolejeunea securifolia* (Nees) Stephani ex Watts ssp. *securifolia* by Thiers & Gradstein.

***Lembidium dendroides Carrington & Pearson* (1888: 1047)**

PROTO: 'On wet earth, Mermaids Glen, Blackheath, Blue Mountains, 3,000ft. September 5, 1885.—*T. Whitelegge*.'

LABEL: NEW SOUTH WALES: Mermaids Glen, Blackheath, Blue Mts, on wet earth, [*T. Whitelegge*] 28, 5 Sep 1885 [2 similarly labelled specimens].

TYPES: BM, NSW (H170 & H171).

COMMENT: NSW specimens annotated by Whitelegge: '*Lembidium dendroides* n. sp., No. 16 P.' and is probably no. 16 of a batch sent to Pearson. Although no collector's number is cited in the protologue all other label data are consistent with these NSW specimens being part of the type. = *Kurzia dendroides* (Carrington & Pearson) Grolle fide Grolle (1963: 175). Still known only from the type (Schuster 1980: 394).

Lepidozia appressifolia *Stephani* (1909: 583)

PROTO: 'Australia orientalis (Watts); Tasmania (Moore)'

LABEL: NEW SOUTH WALES: Katoomba Falls, Blue Mountains, *W.W. Watts 571*, 15 Jan 1902.

TYPES: G, NSW.

COMMENT: NSW specimen of *Watts 571* is labelled 'TYPE' in Watts' handwriting.**Lepidozia buffalona** *Stephani* (Stephani & Watts 1914: 111)PROTO: 'Australia, (Buffalo Creek, near Gladesville; Blue Mountains and Cambewarra): *Watts 977, 941, 978, 965, 982, 914* etc.'LABEL: NEW SOUTH WALES: 1. Centennial Glen, B[lack]heath, *W.W. W[at]ts 1045*, 11 Jan 1911; 2. Horseshoe Falls, Blackheath, wet cliffs, *W.W. W[at]ts 1025*, 7 Jan 1911; 3. Rotunda, Blackheath, *W.W. Watts 1011*, 4 Jan 1911 [3 packets]; 4. Cambewarra Mtn, damp places under cliffs above Baldys, *W.W. W[at]ts 914*, Oct 1907.

TYPES: G, NSW.

COMMENT: *Icones Hep. Steph. 5076* is based on *Watts 977* (Geissler & Bischler 1985: 21).**Lepidozia communis** *Stephani* (Stephani & Watts 1914: 111)PROTO: 'Australia, (Grand Canyon, Blackheath): *Watts, 1014*.'LABEL: NEW SOUTH WALES: Grand Canyon, Blackheath, *W.W. Watts 1014*, 4 Jan 1911 [3 packets in folder].

TYPES: G, NSW.

COMMENT: Name in current use.

Lepidozia crassitexta *Stephani* (Stephani & Watts 1914: 111)PROTO: 'Australia, (Rodriguez Pass, Blackheath): *Watts, 1005*.'LABEL: NEW SOUTH WALES: Rodriguez Pass, Blackheath, *W.W. Watts 1005*, 4 Jan 1911 [3 packets in folder].

TYPES: G, NSW.

COMMENT: Name in current use.

Lepidozia furcatifolia *Stephani* (Stephani & Watts 1914: 112)PROTO: 'Australia, (Horse Shoe Falls, Blackheath): *Watts, 1027*.'LABEL: NEW SOUTH WALES: 1. Horse Shoe [Horseshoe] Falls, Blackheath, wet cliff, *W.W. Watts 1027*, Jan 1911; 2. Horse Shoe [Horseshoe] Falls, Blackheath, wet cliff, *W.W. Watts 1027b*, Jan 1911 [2 packets].TYPES: G, NSW (*Watts 1027*, also? *1027b*).COMMENT: The NSW specimen of *W.W. Watts 1027(b)*, which is labelled 'TYPE' (probably Watts' handwriting), may also be part of the type.**Lepidozia gracillima** *Carrington & Pearson* (1888: 1045)

PROTO: 'on wet rocks, Botany Bay, June, 1885'

LABEL: NEW SOUTH WALES: 1. Botany Bay, on wet rocks, [*T. Whitelegge 24*], June 1885; 2. Botany Bay, on wet rocks, [*T. Whitelegge 24*], June 1885.

TYPES: BM or MANCH?, NSW.

COMMENT: The former NSW specimen has been annotated by Whitelegge: '*Lepidozia gracillima* n. sp. pLXXVII, 51 TW. W.H.P.' and the latter: '24 N. sp. No 51.P. *Lepidozia gracillima*'. Probably no. 51 of a batch sent to Pearson by Whitelegge.

Lepidozia hastatistipula Stephani (Stephani & Watts 1914: 113)

PROTO: 'Australia, (Healesville, Vict.): Watts, 966.'

LABEL: VICTORIA: 'Airlee', Healesville, (by summer house), W.W. Watts 966, 22 Dec 1906 [2 similarly labelled packets in specimen folder].

TYPES: G, NSW (H159).

COMMENT: Name in current use.

Lepidozia lateconica Stephani (Stephani & Watts 1914: 113)

PROTO: 'Australia, (Barron Falls, North Queensland; leg Mrs Brotherton) Watts, 976.'

LABEL: QUEENSLAND: 1. Kuranda, Mrs Brotherton, W.W. Watts 976, June 1910 [3 similarly labelled packets in folder]; 2. Kuranda, Barron Falls, Mrs Brotherton, W.W. Watts 976, June 1910 (NSW H160).

TYPES: G, NSW.

COMMENT: = *Kurzia lateconica* (Stephani) Grolle fide Grolle (1963: 175). Apparently known only from the type (Schuster 1980: 387).

Lepidozia microstipula Stephani (Stephani & Watts 1914: 114)

PROTO: 'Australia, (Lane Cove River, above Fig Tree Bridge): Watts, 958.'

LABEL: NEW SOUTH WALES: 1. Lane Cove River, under rocks, old residential area, first wharf above Fig Tree, W.W. Watts 958, Nov 1910; 2. Near Lane Cove River, above Hunters Hill, under rock in old residential area, W.W. Watts 958, Nov 1910 [2 packets in folder: NSW H161].

TYPES: G, NSW.

COMMENT: Name in current use.

Lepidozia multifida Stephani (Stephani & Watts 1914: 114).

PROTO: 'Australia, New South Wales (Blue Mountains): Watts, 974, 975, 964.'

LABEL: NEW SOUTH WALES: 1. Water Nymphs Dell, Wentworth Falls, W.W. Watts 974, 3 Jan 1910; 2. Nymphs Dell, Wentworth Falls, W.W. Watts 975, 3 Jan 1910 [3 packets in folder, 2 mistranscribed as Jan 1911]; 3. Loc?, [W.W. Watts] 964, 1910.

TYPES: G, NSW.

COMMENT: Name in current use.

Lepidozia nova Stephani (Stephani & Watts 1914: 115)

PROTO: 'Australia, (Blue Mountains and Mount Wilson): Watts, 1033, 1054 and 1066.'

LABEL: NEW SOUTH WALES: 1. Mermaids Glen, Blackheath, W.W. Watts 1054, 3 Jan 1911; 2. Mermaids Glen, Blackheath, W.W. [Watts] 1054, 3 Jan 1911 (NSW H162); 3. Track from Horse Shoe [Horseshoe] Falls to Jungle, Blackheath, W.W. Watts 1033, Jan 1911; 4. Horse Shoe [Horseshoe] Falls to jungle, Blackheath, W.W. [Watts] 1033, 7 Jan 1911 (NSW H163); 5. 'Yarrawa', Mt Wilson, W.W. Watts 1066, 11 Jan 1911 [2 packets].

TYPES: G, NSW.

COMMENT: Name in current use.

Lepidozia quadriseta Stephani (1909: 582)

PROTO: 'N. South Wales (Watts).'

LABEL: NEW SOUTH WALES: 1. Horseshoe Falls, Blackheath, W.W. *Watts* 1027, 7 Jan 1911 (NSW H164); 2. E Ballina, swamp, heath, W.W. *Watts* 433, 11 June 1902.

TYPES: G, NSW.

COMMENT: NSW H164 has been annotated 'type' by Watts. = *Telaranea tetradactyla* (Hook. f. & Taylor) E. Hodgson fide Grolle (1963).

Lepidozia quadristipula Stephani (Stephani & Watts 1914: 115)

PROTO: 'Australia, (Rotunda, Neate's Glen, Blackheath): Watts 1009.'

LABEL: NEW SOUTH WALES: 1. Rotunda, Blackheath, W.W. *Watts* 1009, 4 Jan 1911; 2. Rotunda, Blackheath, W.W. *Watts* 1009, 4 Jan 1911 (NSW H167).

TYPES: G, NSW.

COMMENT: Name in current use.

Lepidozia septemfida Stephani (1909: 588)

PROTO: 'Australia, N.S. Wales (Watts).'

LABEL: NEW SOUTH WALES: Katoomba Falls, W.W. *Watts* 574, 15 Jan 1902.

TYPES: G, NSW (H212).

COMMENT: NSW specimen annotated by Watts: 'Co-type. Mixed with *L. appressifolia*.' Name in current use.

Lepidozia tripilosa Stephani (Stephani & Watts 1914: 116)

PROTO: 'Australia, (Centennial Glen, Blackheath): Watts, 1043.'

LABEL: NEW SOUTH WALES: Centennial Glen, Blackheath, W.W. *Watts* 1043, 10 Jan 1911 [2 similarly labelled packets in specimen folder].

TYPES: G, NSW.

COMMENT: Icones Hep. Steph. n. 5100 is based on *Watts* 1043 (Geissler & Bischler 1985: 47).

Leptocolea wattsiana Stephani (1916: 856)

PROTO: 'Australia. New South Wales.'

LABEL: NEW SOUTH WALES: 1. nr. beach, H. C[ree]k, [W.W. *Watts*] 232, Mar 1901; 2. on tree by J.W. Sharpe's sliprails, nr. beach, above Ballina, W.W. *Watts* 232, Mar 1901 [both specimens in same folder].

TYPES: G, NSW.

COMMENT: Specimen 1 above further annotated by Watts: '*Cololejeunea wattsiana* sp. n. in letter from Dr. Levier July '02'. It has not been possible to identify the locality of 'H. Ck' or 'J.W. Sharpe's sliprails'. Specimen of *Watts* 232 at G cited as holotype and NSW specimen as isotype by Thiers (1988: 123) who does not indicate if the G specimen has been labelled as the type by Stephani. Reduced to synonymy under *Cololejeunea lanciloba* Stephani by Thiers.

Marchantia paludicola Stephani (Stephani & Watts 1914: 120)

PROTO: 'Australia, New South Wales, (Cambewarra): Watts, 1106, 1102.'

LABEL: NEW SOUTH WALES: 1. above Baldys, Cambewarra Mountain, W.W. W[atss] 1106, 18 Oct 1907; 2. above Baldys, Cambewarra Mountain, [W.W.] Watts 1106, Oct 1907; 3. In swamp, 3900 ft [1170 m], Mt Werong, R.H. Cambage, 4 Dec 1911, Hb Watts No. 1102.

TYPES: G, NSW.

COMMENT: NSW specimen collected by R.H. Cambage has been further annotated 'co-type' whilst the other two NSW specimens have been annotated 'type' by Watts. = *Marchantia berteriana* Lehm. & Lindenb. fide Bischler-Cause (1989: 82). NSW specimens cited as syntypes by Bischler-Cause.

Marchantia wattsiana Stephani ex Bonner (Bonner 1953: 112)

PROTO: 'Australia: New South Wales, on stone in Whian Creek, Federal, Richmond River, 13 Aug 1902, Watts, s.n. (Hb Levier 4529), lectotype; Miami [= Minmi] July 1901 Watts 317 (Hb Levier 3116 bis); Richmond River, 1901, Watts 308 (Hb Levier 3111).'

LABEL: NEW SOUTH WALES: 1. Whian Whian Creek, Federal, Richmond River (above Crossing), on stone, W.W. Watts 512, 13 Aug 1902; 2. Richmond River, probably Shaw's Bay, W.W. Watts 308, 1901; 3. loc.?, perhaps Minmi, W.W. Watts 317, July 1901.

TYPES: G (ex Hb Levier 4529: lectotype), NSW.

COMMENT: In 1953 Bonner published a Stephani manuscript, adding to the protologue a lectotype and list of specimens based on Stephani's herbarium at G; he did not examine Watts' collections held at NSW. = *Marchantia pileata* Mitten fide Bischler-Cause (1989: 119) who considers Stephani's description to be based, at least partly, on Watts 308.

Mastigobryum corbieri Stephani (Stephani & Watts 1914: 122)

PROTO: 'Australia, (Kingwell, Wyong): Watts, 971^b.'

LABEL: NEW SOUTH WALES: gully south of house, 'Kingwell', Wyong, W.W. Watts 971^b, Nov 1907 [2 similarly labelled specimens].

TYPES: G, NSW (H286 & H301).

COMMENT: *Mastigobryum* is an illegitimate name but this taxon has never been formally transferred to *Bazzania*, where it belongs (Scott & Bradshaw 1986: 67). *Watts 971a* is the type specimen of *Balantiopsis subkingwella*, see above.

Mastigobryum dentistipulum Stephani (Stephani & Watts 1914: 122)

PROTO: 'Australia, New South Wales (Lane Cove River and Valley of Waters): Watts 1096, 1129, 1104.'

LABEL: NEW SOUTH WALES: 1. Lane Cove River, W.W. Watts 1096, May 1912 (NSW H287); 2. Valley of Waters [Wentworth Falls], W.W. Watts 1129, [3 packets in folder: NSW H288]; 3. Lane Cove River, W.W. Watts 1104, May 1912.

TYPES: G, NSW.

COMMENT: NSW H287 & H288 have been annotated 'co-type' and 'type' respectively by Watts. *Mastigobryum* is an illegitimate name but this taxon has never been formally transferred to *Bazzania*, where it belongs (Scott & Bradshaw 1986: 67).

Mastigobryum erosifolium *Stephani* (Stephani & Watts 1914: 122)

PROTO: 'Australia (Cambewarra Mountain): Watts legit, 917.'

LABEL: NEW SOUTH WALES: 1. Damp places under cliff, Cambewarra Mountain, W.W. *Watts* 917, Oct 1907; 2. Under cliff, above Baldy's, Cambewarra, W.W. *Watts* 917, Oct 1907.

TYPES: G (15074), NSW (H302 & H289).

COMMENT: = *Acromastigum colensoanum* (Mitten) Evans ex Reimers fide Grolle (1972: 547).

Mastigobryum gracillimum *Stephani* (Stephani & Watts 1914: 123)

PROTO: 'Australia, New South Wales, (Wyong and Valley of Waters): Watts, 939, 987.'

LABEL: NEW SOUTH WALES: 1. Gully south of house, 'Kingwell', Wyong, W.W. *Watts* 939, Nov 1907 [2 specimens in separate folders: original Watts packet is NSW H291]; 2. S Wyong, W.W. *Watts* 987, 1909 (NSW H292); 3. Valley of Waters [Wentworth Falls], W.W. [*Watts*] 1130, Sep 1912 (NSW H290) [3 packets in folder].

TYPES: G, NSW.

COMMENT: The NSW specimens of *Watts* 1130 are annotated: '*Mastigobryum gracillimum* St. sp. n. Jan 1913' (the specimens of *Watts* 939 & 987 are similarly labelled but lack the 'sp. n.' designation). *Mastigobryum* is an illegitimate name but this taxon has never been formally transferred to *Bazzania*, where it belongs (Scott & Bradshaw 1986: 67).

Mastigolejeunea wattsiana *Stephani* (1912: 780)

PROTO: no specimens cited.

LABEL: NEW SOUTH WALES: Wallace's Hill, Possum Shoot, R[ichmond] R[iver], on vine, W.W. [*Watts*] 350, Oct 1901.

TYPES: FH, G (21835), NSW.

COMMENT: G 21835 (*Watts* 350) cited as holotype and FH as isotype by Thiers & Gradstein (1989: 50) who do not indicate if the G specimen has been labelled as the type by Stephani. Reduced to synonymy under *Mastigolejeunea ligulata* (Lehm. & Lindenb.) Schiffner by Thiers & Gradstein.

Metzgeria howeana *Stephani* (Stephani & Watts 1914: 126)

PROTO: 'Australia, Mount Wilson and Wyong: Watts legit 1065.'

LABEL: NEW SOUTH WALES: 1. 'Yarrowah', Mt Wilson, W.W. [*Watts*] 1065, 11 Jan 1911; 2. gully over mountain SW of 'Kingwell', Wyong, W.W. *Watts* 948, 24 Aug 1910.

TYPES: G, NSW.

COMMENT: There are four other NSW collections from Yarrowah (*Watts* 1058, 1061, 1069 & 1072), all dated 11 Jan 1911 and annotated 'co-type'. The NSW specimen from Wyong is annotated 'type' in Watt's hand. This species is not listed in Scott & Bradshaw (1986); name in current use.

Pallavacinia campanulata *Stephani* (Stephani & Watts 1914: 127), as '*Pallavacinius*'

PROTO: 'Australia, New South Wales, (Valley of Waters): Watts legit, 1111.'

LABEL: NEW SOUTH WALES: Valley of Waters [Wentworth Falls], [W.W.] *Watts* 1111, Sep 1912 [2 packets in folder].

TYPES: G, NSW (H224).

COMMENT: Name in current use.

***Physocolea casuarinae Stephani* (1916: 909)**

PROTO: 'Australia orientalis. New South Wales.'

LABEL: NEW SOUTH WALES: 1. Fig Tree, Lane Cove R[iver], [Sydney], on *Casuarina*, W.W. W[atts] 792, 1903; 2. bottom of gully above bridge, Fig Tree, Lane Cove River, on *Casuarina*, [W.W.] W[atts] 792, 1903 [both specimens are in the same folder].

TYPES: G (17524), NSW.

COMMENT: G 17524 (Watts 792) cited as holotype by Thiers (1988: 134) who does not indicate if the G specimen is labelled as the type by Stephani. Reduced to synonymy under *Cololejeunea minutissima* (Smith) Schiffner by Thiers.

***Plagiochasma queenslandicum Stephani* (1917: 9)**

PROTO: 'Australia, Queensland (Simmonds legit).'

LABEL: QUEENSLAND: ex Hb Simmonds, W[atts] Q258, Aug 1887.

TYPES: G, NSW.

COMMENT: = *Reboulia hemisphaerica* fide Bischler (1979: 76). Recently reinstated as *Reboulia queenslandica* (Stephani) Hicks fide Hicks (1992: 114) who cites the type as nr Brisbane, J.H. Simmonds 16340, 1887, Herb. W[atts] 611 (4490) (G, FI [FH]).

***Plagiochila rossii Stephani* (Stephani & W[atts] 1914: 130)**

PROTO: 'Australia, N.S. Wales (The Jungle, Blackheath): W[atts] 1035.'

LABEL: NEW SOUTH WALES: Blackheath, The Jungle, W.W. W[atts] 1035, 7 Jan 1911 [3 packets in folder].

TYPES: G, NSW (H344).

COMMENT: = *Plagiochila baileyana* Stephani fide Inoue (1986: 377).

***Plagiochila serrifolia Stephani* (Stephani & W[atts] 1914: 130)**

PROTO: 'Australia, New South Wales (Wyong): W[atts] 1100.'

LABEL: NEW SOUTH WALES: gully over mountain SW of 'Kingwell', Wyong, [W.W.] W[atts] 1100, 23 Aug 1910 [2 packets in folder].

TYPES: G, NSW (H345).

COMMENT: = *Plagiochila vitiensis* Mitten fide Inoue (1986: 364).

***Plagiochila victoriae Stephani* (Stephani & W[atts] 1914: 131)**

PROTO: 'Australia, Victoria (Lorne: leg. Miss E.L. W[atts]) Hb W[atts] 910.'

LABEL: VICTORIA: Lorne, leg. Ethel W[atts], Hb W.W. W[atts] 910, Dec 1909.

TYPES: G, NSW (H370).

COMMENT: = *Plagiochila baileyana* fide Inoue (1986: 377).

Plagiochila watsii Stephani (in Rodway 1918: 105)

PROTO: 'On Fagus log, Pioneer track, Blue Tier (Weymouth). Determined by Stephani.'

LABEL: NEW SOUTH WALES: Pearce's Creek, Richmond River, on tree, *W.W. Watts 19*, 19 June 1900.

TYPES: G (ex Hb Levier 2306), NSW.

COMMENT: Rodway (1918: 105) reported this species from Tasmania, attributing the species description and identification to Stephani. In 1921 Stephani redescribed *Plagiochila watsii* as a new species (the protologue cites the type specimen as: 'Australia. New South Wales (Watts legit)'). The Stephani description does not match the one published in Rodway. Inoue & Schuster (1971) were unable to locate Rodway's specimen, and they suggested that it is probably a species of *Lophocolea*. The G specimen of *Watts 19* is cited as the type by Inoue & Schuster (1971: 221) and Inoue (1986: 371). Further work to clarify the status of this taxon is required!

Porella crawfordii Stephani (1889: 270), as 'cranfordi'

PROTO: 'New England, Moona near Walcha leg. A. R. Cranford 1884....Queensland, Toowoomba, Hartmann 1876, Queensland F. M. Bailey (Herb. Brotherus Nr. 11). N. S. Wales Rev. Dr. Woolls. 1884'

LABEL: NEW SOUTH WALES: Moona near Walcha, New England, *A.R. Crawford*, 1884.

TYPE: G (lectotype), NSW (H259: isolectotype).

COMMENT: Lectotypified by Hattori (1985: 53). This species was named after its collector, A.R. Crawford, but Stephani appears to have misread the handwritten label. Stephani's spelling, although original, must count as an orthographic error and was amended by Scott & Bradshaw (1986: 79).

Riccardia babindae Hewson (1970b: 85)

PROTO: 'Holotype-The Boulders, Babinda, N. Qld., Hewson 460, viii.1964 (NSW): ISOTYPE (BRI).'

LABEL: QUEENSLAND: Babinda, The Boulders, *H.J. Hewson H64.79.460*, 26 Aug 1964.

TYPE: NSW (179242: holotype), BRI (isotype).

COMMENT: Name in current use.

Riccardia bliklika Hewson var. bliklika (Hewson 1970b: 83)

PROTO: 'Holotype-Vision Falls, Lake Eacham, Atherton Tableland, N. Qld., on weathering basalt in rainforest, Hewson, 387, 8.1964, (NSW): ISOTYPE (BRI).'

LABEL: QUEENSLAND: Lake Eacham, Vision Falls, *H.J. Hewson H64.68.387*, 20 Aug 1964.

TYPES: NSW (179243: holotype), BRI (isotype).

COMMENT: Name in current use.

Riccardia hypipamensis Hewson (1970b: 97)

PROTO: 'Holotype- The Crater, Mt. Hypipamee, N. Qld, on weathering basalt in rainforest, Hewson, 448, 8.1964 (NSW): ISOTYPE (SYD).'

LABEL: QUEENSLAND: Mt Hypipamee, The Crater, *H.J. Hewson H64.76.448*, 24 Aug 1964.

TYPES: NSW (holotype), SYD (isotype-not located).

COMMENT: Name in current use.

Riccardia macdonaldiana Hewson (1970b: 93)

PROTO: 'Holotype- Mossman Gorge, 3/4 mile beyond Mossman Mission, on soil and roots in rainforest, Hewson 326 [= 362], 8.1964, (NSW): ISOTYPE (BRI).'

LABEL: NORTH QUEENSLAND: about 3/4 mile [1 km] beyond Mossman Mission, Mossman Gorge-Creek, soil and roots, rainforest, granite, H.J. Hewson H64.66.362, 19 Aug 1964.

TYPES: NSW (holotype), BRI (isotype).

COMMENT: In the protologue the number is given as 326, a typographical error for the correct number 362.

Riccardia minima Carrington & Pearson (1888: 1055)

PROTO: 'On the surface of peat, in bogs, Coogee Bay, New South Wales, May, 1885.'

LABEL: NEW SOUTH WALES: Coogee Bay, in flower, [T. Whitelegge 45], May 1885.

TYPES: MANCH (Kk854), BM, NSW (H247).

COMMENT: NSW specimen further annotated by Whitelegge: '45 N. sp. *Riccardia minima* No. 37.' i.e. probably no. 37 of a batch sent to Pearson by Whitelegge. MANCH Kk854 cited as holotype by Hewson (1970b: 84). = *Riccardia lobulata* (Colenso) E. Hodgson fide Brown & Braggins (1989: 43).

Riccia areolata Na-Thalang (1980: 78)

PROTO: 'HOLOTYPE: New South Wales, Prospect Reservoir, Na-Thalang 236, 15.vi.1967 (NSW).'

LABEL: NEW SOUTH WALES: Prospect Reservoir, open grassland at roadside, O. Na-Thalang 236, 15 June 1969.

TYPES: NSW (M11480: holotype), SYD (isotype-not located).

COMMENT: Name in current use.

Riccia asprella Carrington & Pearson (1888: 1039)

PROTO: 'On earth, Head of Double Bay, Sydney, July, 1885.'

LABEL: NEW SOUTH WALES: Head of Double Bay, on earth, [T. Whitelegge 50], June 1885.

TYPES: BM or MANCH?, NSW (H94).

COMMENT: The NSW specimen has been further annotated by Whitelegge: '50 N. sp., *Riccia asprella*, No 53.P.'; probably no. 53 of a batch sent to Pearson. NSW specimen cited as holotype by Na-Thalang (1980: 84) but the original specimen is almost certainly held at MANCH or BM.

Riccia blackii Na-Thalang (1980: 81)

PROTO: 'HOLOTYPE: Central Australia, Mt Olga. D. Black (Na-Thalang 87), 11.viii[vi].1966 (NSW); ISOTYPE (SYD).'

LABEL: NORTHERN TERRITORY: top of Mt Olga, along creek bank, D. Black, O. Na-Thalang 87, 11 June 1966.

TYPES: NSW (M11478: holotype), SYD (isotype).

COMMENT: Name in current use.

Riccia bullosa Link var. *vesiculosa* Carrington & Pearson (1888: 1058)

PROTO: 'On earth, Parramatta, August, 1885.'

LABEL: NEW SOUTH WALES: 1. Parramatta, on earth, [T. Whitelegge 49], Aug 1885 (NSW H87); 2. Parramatta North, on earth, [T. Whitelegge 49], 16 Aug 1885; 3. Parramatta (North), on earth, [T. Whitelegge 49], 16 Aug 1885 (NSW H254).

TYPES: BM or MANCH?, NSW.

COMMENT: The NSW specimens are further annotated by Whitelegge: '49, 21 T.W. W.H.P', '49, No. 21, *Riccia bulbosa* var. *vesiculosa*' and 'No. 49 C&P, No. 24W.' respectively. No collector or number are cited in the protologue but the other data and annotations are consistent with the NSW specimens being part of the type. Probably no. 21 of a batch sent to Pearson. Na-Thalang (1980: 116) cites the type as 'Holotype: New South Wales, Parramatta, *Whitelegge 8*, 1885 (NSW H87); isotype (SYD).' No material was located at SYD and the original specimen is almost certainly held at MANCH or BM. = *Riccia vesiculosa* (Carrington & Pearson) Stephani fide Na-Thalang (1980: 114).

Riccia caroliniana Na-Thalang (1980: 72)

PROTO: 'HOLOTYPE: Robin Water Fall, N.T., *Na-Thalang 288 28.i.1969* (NSW); ISOTYPE (SYD).'

LABEL: NORTHERN TERRITORY: c. 2 miles [3 km] from Robin Waterfall, creek bank among grass, O. *Na-Thalang 288 28 Jan 1969*.

TYPES: NSW (holotype; not located), SYD (isotype).

COMMENT: Probably on loan.

Riccia collata Na-Thalang (1980: 122)

PROTO: 'HOLOTYPE: New South Wales, Tunmallallee *Na-Thalang 220 26.iii.1967* (NSW); ISOTYPE (SYD).'

LABEL: NEW SOUTH WALES: Tunmallallee, along creek bank, O. *Na-Thalang 220, 26 Mar 1967*.

TYPES: NSW (M11478: holotype), SYD (isotype).

COMMENT: Name in current use.

Riccia duplex Lorbeer var. *megaspora* Na-Thalang (1980: 128)

PROTO: 'HOLOTYPE: Northern Territory, East Alligator River, R.C. *Carolin 6880 15.v.1968* (NSW).'

LABEL: NORTHERN TERRITORY: margins of lagoon, East Alligator River, about 7 miles [11 km] S Cahills Crossing, R.C. *Carolin 6880, 16 May 1968*.

TYPE: NSW (holotype).

COMMENT: Name in current use.

Riccia longiciliata Na-Thalang (1980: 75)

PROTO: 'HOLOTYPE: Northern Territory, Palm Valley, *Na-Thalang 100A, 15.viii.1966* (NSW); ISOTYPE (SYD).'

LABEL: NORTHERN TERRITORY: Palm Valley, open sandy soil or on pocket of rocks, grow[ing] mix[ed] with *R. albida*, O. *Na-Thalang 100A, 15 Aug 1966*.

YPES: NSW (M11477: holotype), SYD (isotype-not located).

COMMENT: Name in current use.

***Riccia luticola* Na-Thalang (1980: 123)**

PROTO: 'HOLOTYPE: Northern Territory, c. 5 miles south of Katherine. *Na-Thalang* 321, 5.ii.1969 (NSW).'

LABEL: NORTHERN TERRITORY: c. 5 miles [8 km] south of Katherine, margin of swamp, among grass, *O. Na-Thalang* 321, 5 Feb 1969.

TYPE: NSW (2 specimens labelled holo- and isotype respectively).

COMMENT: Name in current use.

***Riccia marginata* Carrington & Pearson (1888: 1056)**

PROTO: 'Parramatta, N.S.W., August 1885.'

LABEL: NEW SOUTH WALES: 1. Parramatta North, on earth, [T. Whitelegge 48], 16 Aug 1885; 2. Parramatta, on earth, [T. Whitelegge 48], Aug 1885.

YPES: BM or MANCH?, NSW (H89 & H91), SYD.

COMMENT: The latter NSW specimen is further annotated by Whitelegge: '48 *Riccia marginata* n. sp., plXXXV, 20 TW WHP.' i.e. probably no. 20 of a batch sent to Pearson. Na-Thalang (1980: 100) cites NSW H89 as the holotype and NSW H91 and SYD as isotypes but there is almost certainly a BM or MANCH specimen which would have been the material seen by Pearson.

***Riccia multifida* (Stephani) Stephani var. *filiformis* Na-Thalang (1980: 131)**

PROTO: 'HOLOTYPE: New South Wales, Coonabarabran, Mallallee Creek, *O. Na-Thalang* 215, 26.iii.1967 (NSW).'

LABEL: NEW SOUTH WALES: along the Mallallee Creek, Coonabarabran, *O. Na-Thalang* 215, 26 Mar 1967 [2 similarly labelled specimens in separate folders].

TYPE: NSW (holotype).

COMMENT: Name in current use.

***Riccia multifida* (Stephani) Stephani var. *torticolla* Na-Thalang (1980: 134)**

PROTO: 'HOLOTYPE: Western Australia, Poison Creek, 50 miles north of Leonora, R.C. *Carolin* 5834, Aug 1966 (NSW).'

LABEL: WESTERN AUSTRALIA: Poison Creek, 50 miles [80 km] N of Leonora, mud at edge of water hole, R.C. *Carolin* 5834, Aug 1967.

TYPE: NSW (holotype).

COMMENT: Name in current use.

***Riccia olgensis* Na-Thalang (1980: 100)**

PROTO: 'HOLOTYPE: Central Australia, Mt Olga, D.R. *Selkirk*, (*Na-Thalang* 136 [= 135]), 20 Aug 1966 (NSW); ISOTYPE (SYD).'

LABEL: NORTHERN TERRITORY: on top of Mt Olga, along creek bank, D.R. *Selkirk* (*O. Na-Thalang* 135), 20 Aug 1966.

YPES: NSW (M11479: holotype), SYD (isotype).

COMMENT: Error in protologue; the number on the NSW and SYD specimens is 135 not 136.

Riccia rorida *Na-Thalang* (1980: 101)

PROTO: 'HOLOTYPE: A.C.T., Cotter R., *Na-Thalang* 191, 27.i.1969 (NSW).'

LABEL: AUSTRALIAN CAPITAL TERRITORY: Cotter River, open grassland along the river bank, *O. Na-Thalang* 191, 27 Jan 1969.

TYPE: NSW (holotype).

COMMENT: Name in current use.

Riccia spongiosula *Na-Thalang* (1980: 113)

PROTO: 'HOLOTYPE: New South Wales, Jubbal, *W.W. Watts* 144 [= 744], 22.vii.1903 (NSW H97).'

LABEL: NEW SOUTH WALES: 1. near Tubbul, Young, ground among rocks, *W.W. Watts* 744, 22 July 1903; 2. near Tubbul, Young, ground among rocks, (this was probably) opposite Fowlers), [*W.W. Watts*] 744, 22 July 1903.

TYPES: NSW (H97: holotype; 244529: isotype).

COMMENT: Both of the NSW specimens are annotated: '*Riccia spongiosa* Steph. sp. n. ms. 1906' in Watt's hand. *Na-Thalang* appears to have made three typographic errors in the protologue but all the other information available (*Na-Thalang* 1980: 114) suggests that these are the correct specimens.

Symphyogyna interrupta *Carrington & Pearson* (1888: 1053)

PROTO: no specimens cited.

LABEL: NEW SOUTH WALES: 1. Mossmans Bay, [near Sydney], wet rocks, [*T. Whitelegge* 42], 8 Aug 1885; 2. Mossmans Bay, [near Sydney], [*T. Whitelegge* 42], 8 Aug 1885.

TYPES: BM or MANCH?, NSW (2 specimens).

COMMENT: NSW H360 has been annotated by Whitelegge: '*Symphyogyna interrupta* n. sp., pLXXXIII, 55 T.W. W.H.P.' and NSW H361 as 'n. sp., No 55P.' Presumably specimen no. 55 of a batch sent to Pearson. No specimen information is given in the protologue and the type material at BM or MANCH has not been sighted but the information on the packet is consistent with that found on other type specimens sent by Whitelegge to Pearson.

Symphyogyna multiflora *Stephani* (1917: 68)

PROTO: 'Australia. New South Wales (Watts legit).'

LABEL: NEW SOUTH WALES: Valley of Waters [Wentworth Falls], *W.W. Watts* 1127, Sep 1912.

TYPES: G, NSW.

COMMENT: The NSW specimen is further annotated: '*Symphyogyna multiflora* St. sp. nov. det Jan.1913 (type)' in Watts' hand.

Trichocolea minutifolia *Stephani* (Stephani & Watts 1914: 134)

PROTO: 'Australia (Wyong): Watts legit, 950.'

LABEL: NEW SOUTH WALES: gully over mountain, SW of 'Kingwell', Wyong, *W.W. Watts* 950, 23 Aug 1910 [4 packets in one folder].

TYPES: G, NSW (H80).

COMMENT: Name in current use.

Trichocolea watsiana Stephani (Stephani & Watts 1914: 135)

PROTO: 'Australia (Wyong): Watts, 986.'

LABEL: NEW SOUTH WALES: 1. Gully beyond mountain SW of 'Kingwell', Wyong, W.W. *Watts* 946, Aug 1910; 2. Gully over mountain SW of 'Kingwell', Wyong, W.W. *Watts* 946, 23 Aug 1910 [3 packets in one folder].

TYPES: G, NSW (H75 & H257).

COMMENT: *Watts* 986 in the protologue is probably a typographic error for 946. The former NSW specimen (H257) has been further annotated: '*Trichocolea watsiana* St. sp. n. 1913, Type' in Watts' hand.

Acknowledgements

P. Geissler kindly checked data sent to Geneva against specimens examined by Stephani in preparing descriptions, to assist with designation of status of types held in Sydney. H.J. Hewson checked data based on her studies.

References

- Bastow, R.A. (1888) Tasmanian Hepaticae. *Pap. & Proc. Roy. Soc. Tasmania* for 1887: 209–289.
- Bastow, R.A. (1914) Victorian hepaticae. *Victorian Nat.* 31(5): 74–81.
- Berrie, G.K. (1963) Australian liverworts I. *Haplomitrium intermedium* sp. nov. (Calobryales). *Proc. Linn. Soc. New South Wales* 87: 191–195.
- Bischler, H. (1979) *Phagiochasma* Lehm. et Lindenb. III. Les taxa d'Asie et d'Océanie. *J. Hattori Bot. Lab.* 45: 25–79.
- Bischler–Causse, H. (1989) *Marchantia* L. The Asiatic and Oceanic taxa. *Bryophytorum Bibliotheca* 38: 1–317.
- Bonner, C.E.B. (1953) De Hepaticis II. An unpublished section of volume 6 of Stephani's *Species Hepaticarum*: the genus *Marchantia*. *Candollea* 14: 101–12.
- Bonner, C.E.B. (1965) *Index Hepaticarum*. Part V. *Delavayella* to *Geothallus*. (Cramer: Weinheim)
- Bonner, C.E.B. (1976) *Index Hepaticarum*. Part VIII. *Jungernannia*. (Cramer: Vaduz)
- Brown, E.A. & Braggins, J.E. (1989) A revision of the genus *Riccardia* S.F. Gray in New Zealand with notes on the genus *Aneura* Dum. *J. Hattori Bot. Lab.* 66: 1–132.
- Carrington, B. & Pearson, W.H. (1888) List of Hepaticae collected by Mr Thomas Whitelegge in New South Wales, 1884–5. *Proc. Linn. Soc. New South Wales* ser. 2, 2(4): 1035–1060.
- Downing, A.M., Ramsay, H.P. & Schofield, W.B. (1991) Mosses in the vicinity of Jenoloan Caves. *Cunninghamia* 2: 371–384.
- Engel, J.J. (1968) A taxonomic monograph of the genus *Balantiopsis* (Hepaticae). *Nova Hedwigia* 16 (1–2): 83–130.
- Geissler, P. & Bischler, H. (1985) *Index Hepaticarum*. Vol. 10. *Lembidium* to *Mytilopsis*. (Cramer: Vaduz)
- Grolle, R. (1963) Ueber *Kurzia* v. Martens. *Rev. Bryol. Lichenol.* 32: 166–180.
- Grolle, R. (1972) Miscellanea hepaticologica 121–130. *J. Hattori Bot. Lab.* 36: 547–551.
- Grolle, R. (1979) Miscellanea hepaticologica 171–180. *J. Bryol.* 10(3): 163–72.
- Hatcher, R.E. (1960) A monograph of the genus *Isotachis*. *Nova Hedwigia* 2(4): 573–608.
- Hattori, S. (1979a) A revision of the Australasian species of the genus *Frullania*, Hepaticae, I. *J. Hattori Bot. Lab.* 45: 323–63.
- Hattori, S. (1979b) A revision of the Australasian species of the genus *Frullania*, Hepaticae, II. *J. Hattori Bot. Lab.* 46: 119–53.
- Hattori, S. (1983) A revision of Australasian species of *Frullania*, Hepaticae. III. *J. Hattori Bot. Lab.* 54: 133–182.
- Hattori, S. (1985) Two Australian species of the genus *Porella* L. (Hepaticae). *Bull. Natl Sci. Mus., Tokyo, Ser. B.*, 11(2): 51–55.

- Hattori, S. (1987) New or little known species of *Frullania* (Frullaniaceae) from Queensland and New South Wales, Australia. *Mem. New York Bot. Gard.* 45: 544–555.
- Hewson, H.J. (1970a) The family Aneuraceae in Australia and New Guinea: I. The genus *Aneura*. *Proc. Linn. Soc. New South Wales* 94(2): 173–193.
- Hewson, H.J. (1970b) The family Aneuraceae in Australia and New Guinea: II. The genus *Riccardia*. *Proc. Linn. Soc. New South Wales* 95(1): 60–121.
- Hicks, M.L. (1992) Queensland liverworts. *Reboulia* Raddi. *J. Hattori Bot. Lab.* 71: 113–117.
- Inoue, H. (1986) Notes on the Plagiochilaceae XIII. Review of the genus *Plagiochila* (Dum.) Dum. in mainland Australia. *J. Hattori Bot. Lab.* 60: 357–378.
- Inoue, H. & Schuster, R.M. (1971) A monograph of the New Zealand and Tasmanian Plagiochilaceae. *J. Hattori Bot. Lab.* 34: 1–225.
- Na-Thalang, O. (1980) A revision of the genus *Riccia* (Hepaticae) in Australia. *Brunonia* 3: 61–140.
- Ramsay, H.P. (1980) Contributions of Rev. W.W. Watts F.L.S. to Australian botany. *Taxon* 29: 455–469.
- Ramsay, H.P. & Seur, J. (1990) Type specimens of bryophytes in Australian herbaria. Pp. 247–252 in P.S. Short (ed.), *History of systematic botany in Australasia*. (Australian Systematic Botany Society Inc: Collingwood)
- Ramsay, H.P., Seur, J., Wilson, P.G. & Goodwin, T. (1990) Register of type specimens of mosses in Australian Herbaria. General introduction and part 1. Special collections at NSW: Lord Howe Island, Vanuatu (New Hebrides). *Telopea* 3: 571–592.
- Rodway, L. (1918) Tasmanian bryophyta. *Pap. Proc. Roy. Soc. Tasmania* 1916: 51–143.
- Schuster, R.M. (1960) Studies on Hepaticae. II. The new family Chaetophyllopsidaceae. *J. Hattori Bot. Lab.* 23: 68–76.
- Schuster, R.M. (1980) Studies on Hepaticae, LIV–LVIII. *Kurzia* v. Mart. [*Microledipodzia* (Spr.) Joerg.], *Mitt., Druceella* Hodgs. and *Isolembidium* Schust. *J. Hattori Bot. Lab.* 48: 337–421.
- Scott, G.A.M. (1985) *Southern Australian Liverworts*. Australian Flora and Fauna Series 2.
- Scott, G.A.M. & Bradshaw, J.A. (1986) Australian liverworts (Hepaticae): annotated list of binomials and check-list of published species with bibliography. *Brunonia* 8: 1–171.
- Scott, G.A.M. & Pike, D.C. (1988) Revisionary notes on *Fossombronina*. *Bryologist* 91(3): 193–201.
- Stephani, F. (1884) Die Gattung *Radula*. *Hedwigia* 23: 113–6, 129–137, 145–9, 161–3.
- Stephani, F. (1889) Hepaticae Australiae 1–3. *Hedwigia* 28: 128–135, 155–175, 257–278.
- Stephani, F. (1906–09) *Species Hepaticarum* 3 (Georg: Geneva) [Relevant page dates: 517–693 (1909)]
- Stephani, F. (1909–12) *Species Hepaticarum* 4 (Georg: Geneva) [Relevant page dates: 97–448 (1910); 449–752 (1911); 753–824 (1912)]
- Stephani, F. (1912–17) *Species Hepaticarum* 5 (Georg: Geneva) [Relevant page dates: 1–176 (1912); 849–1008 (1916)]
- Stephani, F. (1917–25) *Species Hepaticarum* 6 (Georg: Geneva) [Relevant page dates 1–128 (1917); 129–176 (1918); 177–240 (1921)]
- Stephani, F. & Watts, W.W. (1914) Hepaticae Australes. *J. & Proc. Roy. Soc. New South Wales* 48: 94–135.
- Thiers, B.M. (1988) The Australian species of *Cololejeunea*. *Beiheft Nova Hedwigia* 90: 113–146.
- Thiers, B.M. & Gradstein, R. (1989) Lejeuneaceae (Hepaticae) of Australia. I. Subfamily Ptychanthoideae. *Mem. New York Bot. Garden* 52: 1–82.
- Watts, W.W. (1902) Notes on some Richmond River hepatics. *Proc. Linn. Soc. New South Wales* 26: 215–216.
- Watts, W.W. (1903) Notes on some Australian hepatics. *Proc. Linn. Soc. New South Wales* 27: 493–494.
- Watts, W.W. (1905) Further notes on Australian hepatics. *Proc. Linn. Soc. New South Wales* 29: 558–560.
- Watts, W.W. (1912) *Census of New South Wales hepatics*. Royal Botanic Gardens, Sydney. (Unpublished manuscript)
- Watts, W.W. & Whitelegge, T. (1902) Census Muscorum Australiensium. *Proc. Linn. Soc. New South Wales* suppl. 27: 1–90.
- Watts, W.W. & Whitelegge, T. (1906) Census Muscorum Australiensium. *Proc. Linn. Soc. New South Wales* suppl. 30: 91–163.

Manuscript received 23 July 1991

Manuscript accepted 4 May 1992